

REVIEW

Open Access

Taurine in morning spot urine for the useful assessment of dietary seafood intake in Japanese children and adolescents

Mari Mori*, Hideki Mori, Atsumi Hamada, Yukio Yamori

From 17th International Meeting of Taurine
Fort Lauderdale, FL, USA. 14-19 December 2009

Abstract

Background: Since our previous report on WHO CARDIAC Study data demonstrated that 24-hour urinary (24U) taurine (Tau) excretion was a useful biomarker of seafood (SF) intake and inversely related to the mortality rates of stroke and coronary heart diseases in the world, we determined that SF is important in the risk reduction of lifestyle related-diseases. The amounts of dietary SF intake are so far estimated from a nutritional survey or 24U Tau excretion.

The sodium/potassium ratio of spot urine (SU) and the 24U ratio were reported to be significantly correlated with. Therefore, we presently examined the relationship of Tau excretion in the morning SU with 24U Tau for simplifying the population comparison and the follow-up of SF intake changes in the process of food education program (FEP).

Methods: After informed consent was obtained, 54 children aged 6-11 years (Children) and 193 adolescents aged 13-18 years (Adolescents) participated in collecting precisely 24U and SU of the first urination on the same day and answered the questionnaire including age and height and weight measurements. The urine samples were measured for creatinine (Cre) and Tau, and the association of these between 24U and SU were analyzed.

Results: The success rates of 24U sampling were very high in Children and Adolescents, 96.4% and 82.4%, respectively. From the result of the multiple regression analysis of SU Tau/Cre and weight we obtained formulas for predicting 24U Tau excretion in Children and Adolescents as follows: Children: $24U \text{ Tau} = 16.3 (\text{weight}) + 314.3 (\text{SU Tau/Cre}) - 175.2$; and Adolescents: $24U \text{ Tau} = 20.2 (\text{weight}) + 644.7 (\text{SU Tau/Cre}) - 569.4$.

Conclusions: The present study established the regression equation to estimate 24U Tau excretion from SU Tau/Cre and weight. These formulas are expected to contribute to the estimation of fish and SF intake and the follow-up of the change of the dietary behavior by FEP in Children and Adolescents.

Background

The average life expectancy of Japanese, presently the top of the average of the males and females in the world markedly increased after World War II by the improvements in sanitation as well as in diets. Japanese school lunch system started initially to solve the problem of malnutrition by supplying flour and skimmed milk

resulted later in the increased rate of the intake of meat, eggs, milk and dairy products, and Western style meals became commonly eaten by school children. This accelerated the loss of traditional Japanese dietary custom to eat rice, fish, soybean, vegetable and sea weeds [1,2]. The rapid Westernization of dietary custom increased health problem related to metabolic syndrome [3], such as obesity, hypertension, dyslipidemia and hyperglycaemia, not only in adults but also in children [4-7].

The Basic Law on Dietary Education, ("Shoku-Iku") was enacted in 2005, to promote the well-balanced

* Correspondence: mmori@mukogawa-u.ac.jp
Institute for World Health Development, Mukogawa Women's University,
Nishinomiya, 6638143, Japan
Full list of author information is available at the end of the article

dietary custom by the education for children and their families. Although, many food educational programs (FEP) were introduced to schools and also to local communities [8], no good objective methods were available to check the improvement of the dietary custom in the children who participated in the programs.

In our world-wide epidemiological investigation, "WHO-coordinated CARDIAC (Cardiovascular Diseases and Alimentary Comparison) Study" [9,10], we could estimate the dietary intake of salt, vegetables and fruit, soy bean and sea food (SF) products by analysis of 24-hour urine (24U) excretions of the biomarkers of these diets such as sodium, potassium, isoflavones and taurine (Tau) [11-14].

Therefore, in the present study we tried to develop a new method for estimating food intake by checking morning spot urine (SU) samples instead of 24U samples to confirm the change of the dietary behavior by FEP, particularly the improvement of the intake of SF, important Japanese traditional food components by examining correlation of urinary Tau excretion, a biomarker of SF intake in 24U and the SU samples from the same individuals.

Methods

Study subjects


We conducted many FEP's for children or adolescents since 2002 in Japan. The subjects of the present study were participants of FEP for elementary school children (Children) aged 6-11 years in Awaji-shima in 2002, and for junior and senior high school girls (Adolescents) aged 13-18 years in 2009, at Mukogawa Women's University (MWU) Junior and Senior High School in Nishinomiya, Japan.

Study design

The protocol for collecting 24U and SU is illustrated in Fig.1. The special jar called "Aliquot cup" which we used for WHO CARDIAC Study was adopted to collect 24U by sampling one fortieth of the voided urine each time of urination [15,16]. After filling up the questionnaires including age and height and weight measurements, all participants were given aliquot-cups and instructed to collect precisely 24U by discarding the first voided urine in the morning and then collecting all urine samples up to the last urine to be sampled in the following morning at the same time when the first urine was discarded in the previous morning. The SU sample (10ml) was collected from the last morning urine sample after sampling one fortieth of the voided urine for 24U collection, and after collecting both SU and 24U samples, participants brought their samples to school. The study protocol was approved by the ethical committee of Mukogawa Women's University, and all participants and their parents gave informed consent for the participation before enrollment.

Urine sample analysis

After 24U volume was collected, urine samples were shared into sampling tubes for the measurement of creatinine (Cre) (mg/dl) concentration by standard laboratory methods at SRL Inc. (Tokyo, Japan) and Tau ($\mu\text{mol/l}$) were estimated by high performance liquid chromatography (HPLC) (GL Sciences, Tokyo, Japan). In order to estimate the whole 24h urinary Tau ($\mu\text{mol/day}$) excretion, we multiplied Tau concentration by the urinary volume of 24 hour. To estimate of 24U Tau from Tau/Cre ratio calculated from SU, we measured Cre and Tau concentrations in SU, and developed a



formula to estimate 24U Tau excretion from checking Tau/Cre ratio in SU in Japanese Children and Adolescents.

Statistical analysis

Statistical analysis used Stat view ver. 5.0 for Macintosh. The SU Tau/Cre ratio was determined from their respective Tau and Cre concentrations of SU. Liner forward stepwise regression analysis was employed to calculate regression equation to predict 24U Tau. The correlation between predicted 24U Tau excretion from SU Tau excretions and actual 24U Tau excretion was then assessed. The comparison between estimated and measured values by 24U collection and sex difference in children data were carried out by student's t-test. Data were expressed as the mean \pm SD, if not specified. Values of $p < 0.05$ were considered statistically significant.

Results

Characteristics of the participants in this study

Characteristics of the participants are summarized in Table 1. From among 432 participants of our FEP, 56 Children (25 boys and 27 girls and 364 Adolescents (all females) tried to collect 24U, and Children and Adolescents 54 and 300, respectively could collect 24U successfully. These were very high success rates of 24U sampling in Children and Adolescents, 96.4% and 82.4% respectively. Further analysis was done in 54 Children and 193 Adolescents, who could collect both 24U and SU successfully on the same day before their participation in our FEP.

There were no significant differences between boys and girl in Children, and significant differences were noted between Children and Adolescents in age, weight, height, 24U Tau and 24U Cre (Table 1). Because of these differences, we developed different formulas for

Children and Adolescents to estimate 24U Tau excretion from SU samples.

Relationship in 24U Tau and SU Tau excretion

As shown in Table 2, among correlation coefficients of 24U Tau excretion with personal characteristics and SU variables, the highest positive correlation scores with 24U Tau excretion were variables of SU Tau/Cre in Children ($R^2=0.86$) and in Adolescents ($R^2=0.69$).

Table 3 shows the result of the multiple liner forward stepwise regression analysis. "SU Tau/Cre and weight ($R^2=0.76$)" in Children was lower determination coefficient than "SU Tau/Cre and age ($R^2=0.79$)" and "SU Tau/Cre and height ($R^2=0.78$)" of 24U Tau. But R^2 is equally high, so that weight was commonly selected in both Children and Adolescents. Then we established the regression equation to predict 24U Tau excretion from SU Tau/Cre and weight in Children and in Adolescents. The obtained formulas for predicting 24U Tau excretion in Children and Adolescents were as follows;

Children: $24U \text{ Tau} = 16.3(\text{weight}) + 314.3 (\text{SU Tau/Cre}) - 175.2$,

Adolescents: $24U \text{ Tau} = 20.2(\text{weight}) + 644.7 (\text{SU Tau/Cre}) - 569.4$.

Discussion

In the results of this study, we indicated the validity of a simple method to estimate the 24U Tau excretion by using SU Tau/Cre and weight data. There are several methods for evaluating salt intake, such as dietary recall or records from 24 to 96 hours, food frequency questionnaires or 24U collection. Generally, 24U collection is considered to be the most reliable method to evaluate salt intake [17,18]. However, in this method participants

Table 1 Characteristics of this study participants

	Boys (n=25)	Children Girls (n=29)	Total (n=54)	Adolescents (n=193)
Age (year)	8.2 \pm 1.4	8.0 \pm 1.4	8.1 \pm 1.4	16.7 \pm 1.3 **
Weight (kg)	31.2 \pm 7.2	29.3 \pm 8.6	30.2 \pm 7.9	49.5 \pm 6.9 **
Height (cm)	134.0 \pm 11.0	131.1 \pm 11.1	132.4 \pm 10.1	156.2 \pm 5.5 **
24U volume (ml)	769.3 \pm 271.1	738.2 \pm 289.1	752.6 \pm 278.7	906.1 \pm 582.4
24U Tau ($\mu\text{mol/day}$)	733.4 \pm 790.9	759.4 \pm 905.5	747.3 \pm 847.1	1106.4 \pm 630.6 **
SU Tau ($\mu\text{mol/l}$)	835.3 \pm 433.2	1370.1 \pm 2472.0	1122.5 \pm 1840.0	1494.6 \pm 1085.7
SU Cre (mg/dl)	97.0 \pm 33.1	90.5 \pm 40.0	93.5 \pm 36.8	139.2 \pm 73.9 **
SU Tau/Cre	0.99 \pm 0.80	1.70 \pm 0.30	1.12 \pm 1.23	1.04 \pm 0.69

24U: 24-h urine, SU: morning spot urine, Tau: taurine, Cre: creatinine. **: $p < 0.0001$ vs. total of Children.

Table 2 The correlation coefficients of 24U Tau excretion with variables of personal characteristics and SU

Variable	Children (n=54)		Adolescents (n=193)	
	Coeff.	p value	Coeff.	p value
Age (years)	0.432	0.0013	0.103	0.1559
Weight (kg)	0.243	0.0764	0.153	0.0337
Height (cm)	0.396	0.0028	0.125	0.0835
SU Cre (mg/dl)	-0.232	0.919	-0.010	0.8956
SU Tau (μmol/l)	0.775	<0.0001	0.632	<0.0001
SU Tau/Cre	0.860	<0.0001	0.686	<0.0001

Coeff: correlation coefficients, 24U: 24-h urine, SU: morning spot urine, Tau: taurine, Cre: creatinine.

must carry urine jars for 24 hours. And sometimes participants cannot obtain urine samples precisely because of forgetting the urine collection. There have been reported some studies that evaluated the effectiveness of SU as a simple method instead of 24U collection [19-21].

In our previous "WHO CARDIAC Study", 24U collection by using "Aliquot cups" was proven to be the useful and practical method to evaluate the different dietary habits of 61 populations in 25 countries [11-14]. In this study we reported 24U Tau excretion was positively correlated with SF intake and inversely correlated with the age adjustment mortality of coronary heart disease (CHD) and stroke [22-26]. In addition, fish and n-3 polyunsaturated fatty acid intake were reported to reduce the risk of CHD and sudden cardiac death in Western countries [27-31].

The risk of CHD was 40% lower among Japanese persons at the highest quintile of fish intake 8 times per week (180g/day) than those who ate fish once per week (23g/day) [32]. Moreover, fish intake beneficially affects the development of type 2 diabetes [33,34]. Higher total fish intake was proven to be associated with 25% reduction of diabetes [35].

On the other hand, dietary habits of Japanese children and adolescents have been Westernized dramatically in the past few decades [4], and the Westernized diet and excessive or imbalanced intake of saturated fatty acids may be important in the pathogenesis of common life-style-related diseases [36] and the prevalence of the risks related to metabolic syndrome increases even in preteen Japanese children [37,38].

Eating fish and SF is a recommendable dietary habit for children's health, and morning SU collection to analyze Tau is an easy method to estimate the intake of fish and SF in children. In addition, Tau itself has been reported to be needed for the development of babies, because Tau is contained a lot in mother's milk [39], and there are many reports, that Tau is effective nutrient for the prevention of CVD [25,40], and diabetes [41]. It is important to execute FEP to make children understand good eating habits of eating fish and SF regularly which are Japanese traditional food. We think that checking SU Tau of participants in our FEP is effective for the confirmation of dietary habit improvement and for the promotion of fish and SF intake.

The limitations of the present study are the limited number of participants in Children and that only girls are studied in Adolescence. In order to apply our formulas for 24U Tau estimation widely to Children and Adolescents, we should collect 24U and SU samples from more children and also from male adolescents.

Conclusions

The present study analyzed the association of SU Tau/Cre with 24U Tau to propose formulas to estimate 24U Tau from SU in Children and Adolescents. These formulas if utilized generally are expected to contribute to the estimation of fish and SF intake and the follow-up of the change of the dietary behavior by FEP and further to the confirmation of the health effect of fish and SF intake.

Table 3 Multiple regression analysis of SU Tau/Cre and weight to predict assessment of 24U Tau excretion

	Children (n=54)				Adolescents (n=193)			
	β	p-value	β	p-value	β	p-value	β	p-value
SU-Tau/Cre	0.84	<0.0001	0.81	<0.0001	0.81	<0.0001	0.70	<0.0001
Weight (kg)	0.15	0.03					0.22	<0.0001
Age (year)			0.24	0.00				
Height (cm)					0.21	0.004		
R ²	0.76	<0.0001	0.80	<0.0001	0.78	<0.0001	0.52	<0.0001
Adjusted R ²	0.75	<0.0001	0.79	<0.0001	0.77	<0.0001	0.51	<0.0001

β: regression coefficient, 95%CI: the 95% confidence interval, 24U: 24-h urine, SU: morning spot urine, Tau: taurine, Cre: creatinine.

List of abbreviations used

CARDIAC: Cardiovascular Diseases and Alimentary Comparison; 24U: 24 hour urine; Tau: taurine; SU: spot urine; Cre: creatinine; SF: seafood; HPLC: high performance liquid chromatography; FEP: food educational programs; Children: elementary school children; Adolescents: junior and senior high school girls; MWU: Mukogawa Women's University.

Acknowledgements

This article has been published as part of *Journal of Biomedical Science* Volume 17 Supplement 1, 2010: Proceedings of the 17th International Meeting of Taurine. The full contents of the supplement are available online at <http://www.jbiomedsci.com/supplements/17/S1>.

We thank all participants of our food educational programs for the participation and urine sample collection, and also the staff for their cooperation in the food educational programs. This work has been implemented by a study grant from Hyogo Prefecture Health Promotion Association in 2002, supported partly by Yamada Bee Farm Grant for Honeybee Research from Yamada Apiculture Center, Inc. in 2009 (to MM).

Authors' contributions

MM designed research planning and participated in the food educational programs and analyzed the data for writing the manuscript. AH performed urinary analysis by HPLC, HM coordinated research and collected urine samples, YY was a responsible doctor for supervising human study. All authors read and approved the final manuscript. MM gave final approval of the version to be published.

Competing interests

The authors declare that they have no competing interests.

Published: 24 August 2010

References

1. Nakamura T: The integration of school nutrition program into health promotion and prevention of lifestyle-related diseases in Japan. *Asia Pac J Clin Nutr* 2008, **17** Suppl 1:349-351.
2. Melby M K, Utsugi M, Miyoshi M, Watanabe S: Overview of nutrition reference and dietary recommendations in Japan: application to nutrition policy in Asian countries. *Asia Pac J Clin Nutr* 2008, **17** Suppl 2:394-398.
3. Ikeda J, Nagata H: Dietary and life-style characteristics and physical condition of school children. *Nippon Kosshu Eisei Zasshi*. 1994, **41**:294-310, Japanese.
4. Murata M: Secular trends in growth and changes in eating patterns of Japanese children. *Am J Clin Nutr* 2000, **72**(5 Suppl):1379S-1383S.
5. National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents: The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics* 2004, **114**:555-576.
6. Yoshinaga M, Sameshima K, Tanaka Y, Arata M, Wada A, Takahashi H: Association between the number of cardiovascular risk factors and each risk factor level in elementary school children. *Circ J* 2008, **72**:1594-7.
7. Nakanishi T, Li R, Liu Z, Yi M, Nakagawa Y, Ohzeki T: Sexual dimorphism in relationship of serum leptin and relative weight for the standard in normal-weight, but not in overweight, children as well as adolescents. *Eur J Clin Nutr* 2001, **55**:989-93.
8. Mori M, Xu JW, Mori H, Ling CF, Wei GH, Yamori Y: Comparative studies on 24-hour urinary excretion in Japanese and Chinese adults and children—need for nutritional education. *Adv Exp Med Biol* 2009, **643**:399-405.
9. WHO Collaborating Center on Primary Prevention of Cardiovascular Diseases, Cardiovascular Diseases for Research Unit: WHO CARDIAC (Cardiovascular Diseases and Alimentary Comparison) Study Protocol and Manual of Operations. Geneva: WHO 1986.
10. Yamori Y: Preliminary report of CARDIAC study: Cross-sectional multicenter study on dietary factors of cardiovascular diseases. CARDIAC Study Group. *Clin Exp Hypertens A* 1989, **11**:957-72.
11. Yamori Y, Nara Y, Mizushima S, Mano M, Sawamura M, Kihara M, Horie R: International cooperative study on the relationship between dietary factors and blood pressure: a preliminary report from the Cardiovascular Diseases and Alimentary Comparison (CARDIAC) Study The CARDIAC Cooperative Study Research Group. *Nutr Health* 1992, **8**:77-90.
12. Yamori Y: Worldwide epidemic of obesity: hope for Japanese diets. *Clin Exp Pharmacol Physiol* 2004, **31**(Suppl 2):S2-4.
13. Liu L, Ikeda K, Yamori Y, WHO-CARDIAC Study Group: Inverse relationship between urinary markers of animal protein intake and blood pressure in Chinese: results from the WHO Cardiovascular Diseases and Alimentary Comparison (CARDIAC) Study. *Int J Epidemiol* 2002, **31**:227-33.
14. Yamori Y, Liu L, Mori M, Sagara M, Murakami S, Nara Y, Mizushima S: Taurine as the nutritional factor for the longevity of the Japanese revealed by a world-wide epidemiological survey. *Adv Exp Med Biol* 2009, **643**:13-25.
15. Yamori Y, Nara Y, Kihara M, Mano M, Horie R: Simple method for sampling consecutive 24-hour urine for epidemiological and clinical studies. *Clin Exp Hypertens A* 1984, **6**:1161-7.
16. Nara Y, Kihara M, Mano W, Horie R, Yamori Y: 'Aliquot cups', simple method for collecting consecutive 24-hour urine samples for epidemiological and clinical studies. *Nutritional Prevention of Cardiovascular Disease* Orlando: Academic Press 1984, 211-6.
17. Yamori Y, Kihara M, Fujikawa J, Soh Y, Nara Y, Ohtaka M, Horie R, Tsunematsu T, Note S, Fukase M: Dietary risk factors of stroke and hypertension in Japan – Part 1: Methodological assessment of urinalysis for dietary salt and protein intakes. *Jpn Circ J* 1982, **46**:933-8.
18. Luft FC, Miller JZ, Lyle RM, Melby CL, Fineberg NS, McCarron DA, Weinberger MH, Morris CD: The effect of dietary interventions to reduce blood pressure in normal humans. *J Am Coll Nutr* 1989, **8**:495-503.
19. Kawasaki T, Ueno M, Uezono K, Kawazoe N, Nakamura S, Ueda K, Omae T: Average urinary excretion of sodium in 24 hours can be estimated from a spot-urine specimen. *Jpn Circ J* 1982, **46**:948-53.
20. Tanaka T, Okamura T, Miura K, Kadowaki T, Ueshima H, Nakagawa H, Hashimoto T: A simple method to estimate populational 24-h urinary sodium and potassium excretion using a casual urine specimen. *J Hum Hypertens*. 2002, **16**:97-103.
21. Mann S J, Gerber L M: Estimation of 24-hour sodium excretion from spot urine samples. *J Clin Hypertens (Greenwich)* 2010, **12**:174-180.
22. Yamori Y, Nara Y, Mizushima S, Sawamura M, Horie R: Nutritional factors for stroke and major cardiovascular diseases: International epidemiological comparison of dietary prevention. *Health Rep* 1994, **6**:22-7.
23. Yamori Y, Liu L, Mori M, Sagara M, Murakami S, Nara Y, Mizushima S: Taurine as the nutritional factor for the longevity of the Japanese revealed by a world-wide epidemiological survey. *Adv Exp Med Biol* 2009, **643**:13-25.
24. Yamori Y, Liu L, Mizushima S, Ikeda K, Nara Y, CARDIAC Study Group: Male cardiovascular mortality and dietary markers in 25 population samples of 16 countries. *J Hypertens* 2006, **24**:1499-505.
25. Yamori Y, Murakami S, Ikeda K, Nara Y: Fish and lifestyle-related disease prevention: experimental and epidemiological evidence for anti-atherogenic potential of taurine. *Clin Exp Pharmacol Physiol* 2004, **31**(Suppl 2):S20-S23.
26. Yamori Y, Liu L, Ikeda K, Miura A, Mizushima S, Miki T, Nara Y, WHO-CARDIAC Study Group: Distribution of twenty-four hour urinary taurine excretion and association with ischemic heart disease mortality in 24 populations of 16 countries: Results from the WHO-CARDIAC study. *Hypertens Res*. 2001, **24**:453-7.
27. Albert CM, Hennekens CH, O'Donnell CJ, Ajani UA, Carey VJ, Willett WC, Ruskin JN, Manson JE: Fish consumption and the risk of sudden cardiac death. *JAMA* 1998, **279**:23-28.
28. Rissanen T, Voutilainen S, Nyyssonen K, Lakka TA, Salonen JT: Fish oil-derived fatty acids, docosahexaenoic acid and docosapentaenoic acid, and the risk of acute coronary events: the Kuopio ischaemic heart disease risk factor study. *Circulation* 2000, **102**:2677-2679.
29. Hu FB, Bronner L, Willett WC, Stampfer MJ, Rexrode KM, Albert CM, Hunter D, Manson JE: Fish and omega-3 fatty acid intake and risk of coronary heart disease in women. *JAMA* 2002, **287**:1815-1821.
30. Albert CM, Campos H, Stampfer MJ, Ridker PM, Manson JE, Willett WC, Ma J: Blood levels of long-chain n-3 fatty acids and the risk of sudden death. *N Engl J Med* 2002, **346**:1113-1118.
31. Lemaitre RN, King LB, Mozaffarian D, Kuller LH, Tracy RP, Siscovick DS: N-3 polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal

- myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr* 2003, **77**:319-325.
32. Iso H, Kobayashi M, Ishikawa J, Sasaki S, Okada K, Kita Y, Kokubo Y, Tsugane S, The JPHC Study Group: Intake of fish and n3 fatty acids and risk coronary heart diseases among Japanese: The Japan public health centre-based (JPHC) study cohort. *Circulation* 2006, **113**:195-202.
 33. Nkondjock A, Receveur O: Fish -seafood consumption, obesity, and risk of type 2diabetes: an ecological study. *Diabetes Metab* 2003, **29**:635-42.
 34. Martín de Santa Olalla L, Nutr Hosp: N-3 fatty acids in glucose metabolism and insulin sensitivity. *Nutr Hosp* 2009, **24**:113-27.
 35. Patel PS, Sharp SJ, Luben R, Khaw KT, Bingham SA, Wareham N J: Association between type of dietary fish and seafood intake and the risk of incident type 2 diabetes - The European prospective investigation of cancer (EPIC) - Norfolk cohort study. *Diabetes care* 2009, **32**:1857-63.
 36. Dolecek TA: Epidemiological evidence of relationships between dietary polyunsaturated fatty acids and mortality in the multiple risk factor intervention trial. *Proc Soc Exp Biol Med* 1992, **200**:177-182.
 37. Yoshinaga M, Tanaka S, Shimago A, Sameshima K, Nishi J, Nomura Y, Kawano Y, Hashiguchi J, Ichiki T, Shimizu S: Metabolic syndrome in overweight and obese Japanese children. 2005, **13**:1135-1140.
 38. Ishikawa M, Arai S, Takano M, Hamada A, Kunimasa K, Mori M: Taurine's health Influence on Japanese high school girls. *J Biomed Sci* 2010, in this issue.
 39. Chesney RW, Helms RA, Christensen M, Budreau AM, Han X, Sturman JA: The role of taurine in infant nutrition. *Adv Exp Med Biol*. 1998, **442**:463-76.
 40. Xu YJ, Arneja AS, Tappia PS, Dhalla NS: The potential health benefits of taurine in cardiovascular disease. *Exp Clin Cardiol* 2008, **13**:57-65.
 41. Schaffer SW, Azuma J, Mozaffari M: Role of antioxidant activity of taurine in diabetes. *Can J Physiol Pharmacol* 2009, **87**:91-9.

doi:10.1186/1423-0127-17-S1-S43

Cite this article as: Mori et al.: Taurine in morning spot urine for the useful assessment of dietary seafood intake in Japanese children and adolescents. *Journal of Biomedical Science* 2010 **17**(Suppl 1):S43.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

